Spitzer Space Telescope spectral observations of AGB stars in the Fornax dwarf spheroidal galaxy

M. Matsuura,^{1,2*} A. A. Zijlstra,³ J. Bernard-Salas,⁴ J. W. Menzies,⁵ G. C. Sloan,⁴ P. A. Whitelock,^{5,6,7} P. R. Wood,⁸ M.-R. L. Cioni,⁹ M. W. Feast,⁶ E. Lagadec,³ J. Th. van Loon,¹⁰ M. A. T. Groenewegen¹¹ and G. J. Harris²

¹National Astronomical Observatory of Japan, Osawa 2-21-1, Mitaka, Tokyo 181-8588, Japan

²Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT

³School of Physics and Astronomy, University of Manchester, Sackville Street, PO Box 88, Manchester M60 1QD

⁴Astronomy Department, Cornell University, 610 Space Sciences Building, Ithaca, NY 14853-6801, USA

⁷NASSP, Department of Mathematics and Applied Mathematics, University of Cape Town, 7701 Rondebosch, South Africa

⁸Research School of Astronomy & Astrophysics, Mount Stromlo Observatory, Australian National University, Cotter Road, Weston, ACT 2611, Australia

⁹SUPA, School of Physics, University of Edinburgh, IfA, Blackford Hill, Edinburgh EH9 3HJ

¹⁰Instituut voor Sterrenkunde, KU Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium

¹¹Astrophysics Group, School of Physical and Geographical Sciences, Keele University, Staffordshire ST5 5BG

Accepted 2007 September 19. Received 2007 September 18; in original form 2007 August 13

ABSTRACT

We have observed five carbon-rich asymptotic giant branch (AGB) stars in the Fornax dwarf spheroidal (dSph) galaxy, using the Infrared Spectrometer on board the Spitzer Space Telescope. The stars were selected from a near-infrared survey of Fornax and include the three reddest stars, with presumably the highest mass-loss rates, in that galaxy. Such carbon stars probably belong to the intermediate-age population (2–8 Gyr old and metallicity of [Fe/H] \sim -1) of Fornax. The primary aim of this paper is to investigate mass-loss rate, as a function of luminosity and metallicity, by comparing AGB stars in several galaxies with different metallicities. The spectra of three stars are fitted with a radiative transfer model. We find that mass-loss rates of these three stars are $4\text{--}7 \times 10^{-6} \,\text{M}_{\odot} \,\text{yr}^{-1}$. The other two stars have mass-loss rates below $1.3 \times 10^{-6} \,\mathrm{M_{\odot} \, yr^{-1}}$. We find no evidence that these rates depend on metallicity, although we do suggest that the gas-to-dust ratio could be higher than at solar metallicity, in the range 240 to 800. The C_2H_2 bands are stronger at lower metallicity because of the higher C/O ratio. In contrast, the SiC fraction is reduced at low metallicity due to low silicon abundance. The total mass-loss rate from all known carbon-rich AGB stars into the interstellar medium (ISM) of this galaxy is of the order of $2 \times 10^{-5} \,\mathrm{M_{\odot} yr^{-1}}$. This is much lower than that of the dwarf irregular galaxy Wolf Lundmark Melotte (WLM), which has a similar visual luminosity and metallicity. The difference is attributed to the younger stellar population of WLM. The suppressed gas-return rate to the ISM accentuates the difference between the relatively gas-rich dwarf irregular and the gas-poor dSph galaxies. Our study will be useful to constrain gas and dust recycling processes in low-metallicity galaxies.

Key words: stars: AGB and post-AGB – stars: atmospheres – stars: carbon – stars: mass-loss.

1 INTRODUCTION

Stars with low and intermediate initial mass (from 1 to $8 M_{\odot}$) loose their atmospheres towards the end of their lives (1–10 Gyr after their birth). This mass loss is intense; approximately 50–80 per

cent of the stellar mass is lost during the asymptotic giant branch (AGB) phase. Understanding this process and the composition of the gas lost from AGB stars is important for understanding both stellar and galactic evolution. First, the stellar wind removes material from AGB stars, reducing their mass and influencing their evolution. Secondly, AGB stars are among the primary sources of metal-enriched gas and dust grains in the interstellar medium (ISM) of galaxies (Maeder 1992). In particular, carbon-rich dust grains are

⁵South African Astronomical Observatory, PO Box 9, 7935 Observatory, South Africa

⁶Astronomy Department, University of Cape Town, 7701 Rondebosch, South Africa

^{*}E-mail: mikako@optik.mtk.nao.ac.jp

Table	1.	Targets
-------	----	---------

Name	Coordinates (J2000)		2MASS designation ^a	Other names	Ref.
Fornax 13-23	02h38m50s6	-34°40′32″.0	02385056-3440319		
Fornax 12-4	02h39m12s3	-34°32′45″.0	02391232-3432450		
Fornax 3-129	02h39m41s6	-34°35′56″.7	02394160-3435567		
Fornax 4-25	02h40m10s2	-34°33′21″.9	02401016-3433218	V20, C10, DM19	1,2,3
Fornax 6-13	$02^{h}40^{m}31^{s}.2$	-34°28′44″.2	02403123-3428441	DM22	1,3

Notes. Ref. 1 – Demers & Irwin (1987); 2 – Westerlund, Edvardsson & Lundgren (1987); 3 – Demers et al. (2002). ^aSkrutskie et al. (2006).

Table 2. Magnitudes and variability.

Name	DI^{a}		WEL^b		IRSF ^c		$2MASS^d$		
	$\langle V \rangle$	Period	V	V - B	$\langle K \rangle$	$\langle J-K\rangle$	J	Н	K_s
Fornax 13-23					13.650 ± 0.4	3.147	16.106 ± 0.0939	14.525 ± 0.0527	12.879 ± 0.0287
Fornax 12-4					12.823 ± 0.3	3.027	14.722 ± 0.0334	13.262 ± 0.0384	12.120 ± 0.0242
Fornax 3-129					13.907 ± 0.4	3.764	<17.663	15.970 ± 0.2049	14.164 ± 0.0695
Fornax 4-25	18.6	317.2	18.43	2.92	13.221 ± 0.2	1.984	14.063 ± 0.0256	13.122 ± 0.0214	12.545 ± 0.0285
Fornax 6-13					13.378 ± 0.1	1.896	14.745 ± 0.0433	13.689 ± 0.0456	13.072 ± 0.0365

Notes. IRSF is a monitoring program providing average magnitudes, while 2MASS measurements present a single-phase observation.

^aDemers & Irwin (1987). ^b2Westerlund et al. (1987). ^cMenzies et al., (in preparation). ^dSkrutskie et al. (2006).

formed solely in carbon-rich AGB stars (Dwek 1998). Theoretical work has suggested that AGB mass-loss rates depend on metallicity (Bowen & Willson 1991; Willson 2006), because the stellar wind is triggered by radiation pressure on dust grains and the dust is made up of astronomical metals, such as oxygen, carbon, silicon, iron and aluminium. Therefore, a study of extragalactic AGB stars is vital, both because it allows us to study mass loss at low metallicity, and because the distances to the parent galaxies are known, so that stellar parameters can be derived with some precision. A high sensitivity mid-infrared (IR) instrument is required for such a mass-loss study; at present this is provided uniquely by the *Spitzer Space Telescope* (Werner et al. 2004).

The Fornax dwarf spheroidal (dSph) galaxy is the second (after Sagittarius) most luminous dSph galaxy known in the Local Group. The metallicity of red giants has been measured to be as low as $[Fe/H]^1 = -0.7$ and -1.5 with a mean value of -1.0 ± 0.3 (Tolstoy et al. 2001). This value is lower than that found in the Large Magellanic Cloud (LMC) (-0.4; median value of red giants in the bar; Cole et al. 2005) and the Small Magellanic Cloud (SMC) (-0.5 or -0.73; Smith & Lambert 1989; van den Bergh 1999). The metallicity of AGB stars has not yet been determined in these galaxies, except for one AGB star in the SMC, which has been estimated at -1.0 ± 0.35 (de Laverny et al. 2006). This is consistent, within the error, with the values quoted above for red giants. Thus, the Fornax dSph is an ideal target to explore the lower metallicity range. Fornax has globular clusters (GCs); their ages range from 7 to 14 Gyr (Strader et al. 2003). Battaglia et al. (2006) found three distinct periods of star-forming activity in the galaxy. The dominant population, to which the AGB stars are most likely to belong, is 2 to 8 Gyr old. There may have experienced star-forming activity until a few times 10⁸ yr ago, but this young population is small. Fornax is gas poor at the present time; Young (1999) reported a non-detection of HI gas, although Bouchard, Carignan & Staveley-Smith (2006) have detected more distant gas possibly associated with Fornax.

The distance modulus of the Fornax dSph is quoted as: $(m - M)_0 = 20.76 \text{ mag}$ (Demers, Grondin & Kunkel 1990), $(m - M)_0 = 20.70 \pm 0.12 \text{ mag}$ (Saviane, Held & Bertelli 2000) and $(m - M)_0 = 20.74 \pm 0.11 \text{ mag}$ (Gullieuszik et al. 2007). These values are consistent and we adopt 20.76 mag in this paper. The Fornax dSph galaxy is elliptical in shape and the major axis is longer than the minor axis by a factor of at least 2 (Irwin & Hatzidimitriou 1995). The thickness of the galaxy is negligible at its distance. The interstellar reddening towards Fornax is $E(B - V) \approx 0.03$ (Demers, Dallaire & Battinelli 2002), and its effect is negligible in the IR.

Demers & Kunkel (1979) found red giants in the Fornax dSph galaxy and Aaronson & Mould (1980) established that some of these are (AGB) carbon stars. Azzopardi et al. (1999) claim to have detected 104 carbon-rich stars in this galaxy. Because of its low metallicity, relatively close distance and reasonable number of AGB stars, the Fornax dSph is an ideal galaxy to study the influence of metallicity on the evolution of AGB stars.

As follow-up to our *Spitzer* spectroscopic survey of AGB stars in the LMC (Matsuura et al. 2006; Zijlstra et al. 2006) and the SMC (Lagadec et al. 2007a), we have undertaken spectral observations of carbon stars in Fornax, using the Infrared Spectrograph (IRS; Houck et al. 2004) on the *Spitzer Space Telescope*. This study aims to investigate the spectra of the AGB stars and to ascertain the influence of metallicity on AGB mass-loss rates and on the composition of molecules and dust grains, down to a metallicity of one tenth of the solar value.

2 TARGETS

Our target selection is based on a monitoring program with the Infrared Survey Facility (IRSF; Glass & Nagata 2000; Menzies et al. 2002) at Sutherland in South Africa (Menzies et al., in preparation). Table 1 lists the five targets together with their coordinates and crossidentifications. Photometric measurements of the targets from the literature are summarized in Table 2. The variability of these stars will be discussed elsewhere (Menzies et al., in preparation), but all

¹ Here we use the chemical abundance ratio of any element *X*, $[X/H] = \log(X/H) - \log(X/H)_{\odot}$ where $\log(H)=12$ is the abundance of hydrogen by number of atoms. The metallicity of the galaxy is represented by iron abundance.



Figure 1. The absolute *K*-band magnitude (M_K) , J - K colour diagram for the five sources in the Fornax dSph galaxy (large filled circles) observed at mid-IR wavelengths. For comparison a large sample of mid-IR sources in the LMC from MSX (Midcourse Space Experiment; small open circles) and field stars from a small area of the LMC bar (small dots) are also shown – see Zijlstra et al. (2006) and Wood et al. (2007) for details. AGB stars are confined approximately to the region below the line marked 'AGB limit'. Stars above this limit are red supergiants with initial masses above $\sim 8 M_{\odot}$, or perhaps foreground stars. Locations of post-AGB stars (PAGB) and tip of FGB (first giant branch) are indicated. Distance moduli of 20.76 and 18.54 mag have been assumed for Fornax and the LMC, respectively.

of them show large amplitude periodic changes, and are most likely Mira variables. This variability will result in spectral changes if the stars are indeed Mira stars (e.g. Hron et al. 1998; Busso et al. 2007).

Fig. 1 shows an IR colour–magnitude diagram of our targets, where they are compared with LMC stars. Our sample is located on the sequence of LMC AGB stars, although the Fornax stars tend to have fainter M_K magnitudes than do LMC stars with similar J - K colours.

Our three red carbon-rich stars (Fornax 13-23, 12-4 and 3-129) are some of the reddest among anything described in a recent near-IR survey of Fornax (Gullieuszik et al. 2007) and probably have the thickest shells of any AGB stars in that galaxy.

The Fornax dSph galaxy is roughly elliptical in shape, although the ellipticity varies slightly depending on the distance from the galaxy centre. All of our targets are located within an ellipse of 0.4° semimajor axis and ellipticity of 0.3 (see Battaglia et al. 2006 their fig. 1; Fig. 2 in our work). A colour–magnitude diagram of Fornax (Battaglia et al. 2006) shows the 'AGB bump', which marks the beginning of the AGB phase; stars in this bump belong to an intermediate age population. Because of the typical age of AGB stars, it is most likely that our targets also belong to the intermediate age population, which has a metallicity of $[Fe/H] \sim -1$ (Battaglia et al. 2006). Contamination by young (<1 Gyr) carbon stars is possible, but not likely. Such young carbon stars would have high luminosities, close to the AGB luminosity limit, which are not found for any of our sample (Fig. 1). In addition, the population of such young carbon stars must be small.

Tolstoy et al. (2003) found [Fe/H] ~ -1.6 among some red giants from optical high spectral resolution observations, but the majority were between -0.9 and -1.3 according to the medium resolution spectroscopic survey (Tolstoy et al. 2001; Battaglia et al. 2006). The median of the 'current' metallicity of red giants is [Fe/H] \sim -1.0 according to spectroscopic observations of over a hundred stars (Pont et al. 2004). Furthermore, a planetary nebula, which is carbon rich, shows [Fe/H] = -1.13 ± 0.18 (Kniazev et al. 2007). We conclude that [Fe/H] ~ -1.0 is a reasonable assumption for the population we are dealing with.

Two of our targets were identified as carbon stars by Demers et al. (2002). The other three targets were not included in their sample, because they selected carbon stars with 1.4 < J - K < 2.0.

3 OBSERVATIONS AND DATA REDUCTION

Our targets were observed with the IRS on board the *Spitzer Space Telescope*, using the low-resolution mode. Only the short-low (SL) module was used with a spectral coverage from 5 to 15 μ m. This wavelength range is covered by three segments SL2, SL1 and a bonus order. The spectral resolution is in the range from 60 to 127. The program ID number is 20357.

Observations were carried out on 2006 January 27 to 30. The exposure time per star was 1440 s (60 s × 24 cycles) for SL2 (5.2 to 8.7 µm) and 1680 s (60 s × 28 cycles) for SL1 (7.4 to 14.5 µm (officially), but extends up to 15.0 µm). We used a nearby star, IRAS F02375-3443 (02^h39^m35^s23, $-34^{\circ}30'37'_{.2}$), to peak-up on.

The data were processed through the S13.2 and S14.0 version of the *Spitzer* Science Centre's (SSC) pipeline. The reduction started from the DROOP products which are equivalent to the commonly used binary coded decimal (BCD) data, but lack flat-field and stray-crosslight removal (the latter is only important for bright sources). Rogue pixels are first flagged using a campaign mask and then removed using the IRSCLEAN² tool. Different cycles (repetitions) were averaged to improve the signal-to-noise ratio (S/N). Finally, one-dimensional spectra were extracted using a variable extraction window set at 4 pixel at the middle of each order. The calibration was performed by dividing the resultant spectrum by that of the calibration star HR 6348 (extracted in the same way as the target) and multiplying by its template (Cohen, Wheaton & Megeath 2003; Sloan et al., in preparation).

4 DESCRIPTION OF THE SPECTRA

Fig. 3 shows the *Spitzer/IRS* spectra of the five target stars; all are carbon rich. The target selection (long period variability) was independent of spectral classification (i.e. O rich versus C rich), and this suggests that high mass loss is only exhibited by carbon stars in this galaxy. Absorption features at 7.5 and 13.7 μ m are due to C₂H₂. There may be some contribution from HCN in the absorption feature at 7 μ m. The emission at 11.3 μ m is a dust excess from SiC. A dip at 6 μ m is part of the 5 μ m CO fundamental, possibly blended with C₃ absorption bands. CO molecules will be present in all cool AGB stars. The identification of C₃ in the Fornax stars is not certain,

² This tool is available from the SSC website http://ssc.spitzer.caltech.edu



Figure 2. Locations of our target stars within the Fornax dSph galaxy, superposed on the Digitized Sky Survey (DSS) image. GCs within the field are also indicated. Coordinates are in J2000 and one square grid box is $15 \times 10 \text{ arcmin}^2$.

because the wavelength coverage of IRS is not favourable for this band, but the presence of C₃ is known for some Galactic (albeit blue) carbon stars (Yamamura et al. 1997; Jørgensen, Hron & Loidl 2000) and therefore it is likely to be present in these extragalactic stars. The SiC excess at 11.3 μ m is strong in Fornax 13-23 and 3-129 and weak in Fornax 12-4. This feature is marginally detected in Fornax 4-25 and is not obvious in Fornax 6-13. The feature at 8.5 μ m is caused by the band gap.

Fornax 6-13 does not show any obvious feature in its IRS spectrum. However, its near-IR colour, J - K > 1.4, suggests that this star is carbon rich (Cioni et al. 2001), and this is confirmed by near-IR spectra (Groenewegen et al., in preparation). Demers et al. (2002) also spectroscopically identified Fornax 4-25 and 6-13 as carbon stars.

4.1 Band strengths

We measured the equivalent width of the molecular bands and the strength of the SiC dust emission, so as to evaluate the metallicity dependence of these features. The equivalent width of the C2H2 molecular bands and SiC are measured following the method of Zijlstra et al. (2006). Figs 4 and 5 show 7.5 and $13.7 \,\mu m C_2 H_2$ equivalent widths as a function of IR colour [6.4]-[9.3]. The [6.4] and [9.3] values are calculated from the Spitzer/IRS spectra, and the definition of these magnitudes is also given by Zijlstra et al. (2006); the bands being selected so as to avoid major molecular features. [6.4]–[9.3] is a measure of the stellar temperature for blue stars, but of circumstellar excess for red stars. Samples in our Galaxy (Milky Way), the LMC and the SMC are also plotted in the figures (Sloan et al. 2003, 2006; Zijlstra et al. 2006; Lagadec et al. 2007a). As seen in the spectra and Table 3, Fornax 13-23 and 3-129 have a large equivalent width at 7.5 μ m (W(7.5) = 0.2–0.25 μ m) with respect to the IR colour [6.4]–[9.3]. Here W is the acronym of 'equivalent width'. These band strengths are some of the largest found in the range 0.5 < [6.4]-[9.3] < 1.0, within the samples from various galaxies. A high value of W(7.5), with respect to IR colour, is found for stars in Fornax and the SMC, while a low W(7.5) is found in our Galaxy. The equivalent width of 13.7 µm is more affected by spike noise because this molecular feature is weak, and the quality of data in 14.5–15.0 µm region is poorer. Nevertheless, Fig. 5 shows that stars in our Galaxy cover the lowest range of 13.7 µm equivalent width, W(13.7) in the range 0.5 < [6.4]–[9.3] < 1.0.

Fig. 6 shows the ratio of SiC excess with respect to the pseudocontinuum (Zijlstra et al. 2006); larger ratios indicate larger SiC excess. A careful positioning of the 'continuum' would be required to measure the SiC band strength precisely (Thompson et al. 2006), but we argue that this ratio represents a good estimate of the approximate SiC strength. The ratio is plotted against the IR colour [6.4]-[9.3] in Fig. 6. A large SiC excess is found for relatively blue (IR colour 0.0 < [6.4]-[9.3] < 0.5) Galactic stars. Stars in the SMC and Fornax have the lowest SiC excess ratio among any of the samples. Two stars in Fornax show a SiC to continuum ratio of ~0.15 at 0.5 < [6.4]-[9.3] < 0.8, much less than found for Galactic stars with similar colour.

5 MASS-LOSS RATE

Using flux calibrated *Spitzer* spectra together with IR photometric data, we derive mass-loss rates for the carbon-rich AGB stars.

5.1 Modelling the spectral energy distribution

Spectral energy distributions (SEDs) are modelled using the radiative transfer code DUSTY (Ivezić & Elitzur 1997). The quality of these SED fits with the observations is illustrated in Fig. 7.

A spherical geometry is assumed for the circumstellar envelope, and the circumstellar shells are filled with material from the radiatively driven wind, which is moving at a constant velocity. The photospheric spectrum of the central star is calculated with a hydrostatic model including molecular opacities (Loidl, Lançon & Jorgensen 2001; Groenewegen et al. 2007). An effective temperature of 2800 K



Figure 3. Spitzer/IRS spectra of carbon-rich stars in the Fornax dSph galaxy. Spectra are smoothed by a factor of 2.5 on the wavelength grid to improve the quality. The wavelength ranges of the molecular bands are indicated by lines, and possible identifications of molecular bands are shown with dotted lines.



Figure 4. The equivalent width of 7.5 μ m C₂H₂ as a function of IR colour [6.4]–[9.3]. Symbols show the host galaxies, MW representing the Milky Way.



Figure 5. The same as Fig. 4 but for the equivalent width of 13.7 $\mu m\,C_2H_2.$

is adopted. We also tried a 2650 K spectrum which often worked well for stars in the LMC and SMC (Groenewegen et al. 2007), but this lower temperature did not result in a good fit for the Fornax spectra. This may be a consequence of the difference between the

Table 3. Equivalent width of molecular bands ($W_{7.5}$ and $W_{13.7}$) for 7.5 and 13.5 μ m C₂H₂, SiC dust emission strength (above the continuum) with respect to the 'continuum' (R_{SiC}), IR fluxes (*f*) at 6.4 and 9.3 μ m, respectively, and the [6.4]–[9.3] colour; definitions are given by Zijlstra et al. (2006).

Name	W _{7.5} (μm)	R _{SiC}	W _{13.7} (μm)	<i>f</i> _{6.4} (mJy)	<i>f</i> 9.3 (mJy)	[6.4]–[9.3] (mag)
Fornax 13-23	0.25	0.14	0.022	7.2	6.7	0.73
Fornax 12-4	0.08	0.04	0.058	10.8	8.4	0.55
Fornax 3-129	0.22	0.15	0.029	12.7	10.6	0.62
Fornax 4-25	0.15	0.03	0.019	2.9	1.4	0.04
Fornax 6-13	0.11	0.01	0.026	2.1	1.2	0.20



Figure 6. The intensity ratio of SiC excess with respect to the pseudocontinuum is plotted against [6.4]–[9.3]; a ratio of 0.0 indicates no SiC excess (continuum only).

samples from the various galaxies; the LMC and SMC stars have redder J - K colours (Fig. 1). As details of carbon to oxygen ratios are unknown, a C/O ratio of 1.1 is assumed in the hydrostatic model. Although this is typical of values found for Galactic carbon stars, at lower metallicity the C/O ratio may well be higher (Matsuura et al. 2002, 2005). The assumed C/O ratio affects most strongly the depth of molecular bands, and we minimize its impact by concentrating on a relatively featureless wavelength range of the spectra for this particular exercise. Furthermore, the elemental abundances (except for carbon) used by DUSTY modelling are the solar values (Anders & Grevesse 1989), which are also inappropriate for stars in Fornax. In spite of these limitations, we prefer the hydrostatic model spectra over simple blackbodies (BBs). We compared SED fits using BB curves the star and hydrostatic model spectra as a heating input of DUSTY. The simple BB requires a lower optical depth to reproduce the SED. For example, Fornax 13-23 and 3-129 are fitted with $\tau_{0.55}$ = 3.0 and 8.25 with BB input, respectively. However, the models with BB input have difficult in fitting near-IR spectra. The low optical depth required by the BB model is probably a consequence of the lack of molecular absorption, thus the more efficient heating of grains especially at shorter wavelength.

The temperature at the inner radius of the dust shell is assumed to be 1000 K (Groenewegen et al. 2007). Dust grains are a mixture of amorphous carbon (Hanner 1988) and SiC (Pégourié 1988). The fractions which give the best fit are 5 per cent SiC and 95 per cent amorphous carbon. We also attempted to fit the observed SiC bands with 3 and 10 per cent SiC mixtures, but the former does not create any SiC bands at 11 μ m, while the latter produces SiC bands that are



Figure 7. SED and fits with DUSTY radiative transfer model.

too strong to fit the spectra of our targets. The grain size distribution is assumed to be a = 0.0005 to 0.25 µm according to a power-law distribution of the form $n(a) \propto a^{-q}$ with q = 3.5 (Mathis, Rumpl & Nordsieck 1977). A gas-to-dust ratio of 200 is the default of DUSTY. Although we note that the ratio might be higher at lower metallicity (van Loon 2000), no dependence has yet been measured. This ratio is important when the optical depth is converted to the mass-loss rate.

The outer radius of the shell is fixed at 10^3 times the inner radius of the circumstellar shell. Changing the outer radius has minor

effects on the SED (van Loon et al. 2003). The DUSTY model fit produces the terminal outflow velocity and mass-loss rate for a luminosity of $10^4 L_{\odot}$. As we derive the luminosity by scaling the emerging spectrum from DUSTY, we recalculate the dust mass-loss rate accordingly.

The DUSTY model itself has an uncertainty of 30 per cent in massloss rate and expansion velocity, as noted in the DUSTY manual.

5.2 Results of radiative transfer code fitting

Figs 7 and 8 show the DUSTY model fits to the spectra and the parameters used for these fits are listed in Table 4. We use three spectra



Figure 8. SED and fits with DUSTY, but with a linear scale to emphasize the SiC dust feature and quality of the fit.

Table 4. Luminosity and mass-loss rate.

Name	$M_{\rm bol}$	L	$\tau_{0.55}$	М	v	$\dot{M}_{ m d}$
Fornax 13-23	-4.92	7389	4.00	4.2×10^{-6}	20	2.2×10^{-6}
Fornax 12-4	-5.54	13012	2.45	4.5×10^{-6}	25	2.4×10^{-6}
Fornax 3-129	-4.97	7748	8.50	6.9×10^{-6}	16	
Fornax 6-13	-4.69^{a}	5721 ^a	<1	${<}1.3\times10^{-6}$		

Notes. M_{bol} : bolometric luminosity (mag) estimated from DUSTY fitting; L: luminosity (L_☉) estimated from DUSTY fitting; $\tau_{0.55}$: optical depth at 0.55 µm for DUSTY model fitting; \dot{M} : gas mass-loss rate (M_☉ yr⁻¹) estimated from DUSTY fitting; v: terminal velocity (km s⁻¹) estimated from DUSTY fitting; \dot{M}_{d} : gas mass-loss rate estimated from near- and mid-IR colour method, assuming a gas-to-dust ratio of 200 (Lagadec et al. 2007b). The K - [11] colour is used. The values quoted in the table are based on method (2) in their paper.

^aSpline fitting to the SED is used instead of DUSTY.

which show clear SiC excesses to fit the 11.3 µm SiC excess and the general shape of the SED. We attempted to fit the other two spectra with negligible SiC excess (Fornax 4-25 and 6-13) and found that the optical depth at 0.55 μ m was $\tau_{0.55}$ < 1. The Fornax 4-25 and 6-13 spectra seem to be photospheric in origin, with little circumstellar contribution to the IR. Integrating over the JHK photometry and the Spitzer spectra of Fornax 6-13, using spline fitting (Zijlstra et al. 2006), we obtained a luminosity of 5721 L_O; DUSTY fitting gave an upper limit to the mass-loss rate of $< 1.3 \times 10^{-6} \,\mathrm{M_{\odot} \, yr^{-1}}$. The same spline fitting is adopted for Fornax 4-25, using VJHK and Spitzer spectra, resulting in a luminosity of 5621 L_O, but with lower accuracy than for the other targets. An upper limit to the mass-loss rate is $<1.3 \times 10^{-6} \,\mathrm{M_{\odot}} \,\mathrm{yr^{-1}}$. These photometric observations are not simultaneous with the Spitzer measurements, in contrast to the study by Groenewegen et al. (2007), increasing the uncertainty in the instantaneous luminosities.

There are some features which the DUSTY models do not fit well, in particular the 7.5 and 13.7 μ m absorptions due to C₂H₂. A similar problem was also found in fitting some LMC and SMC AGB spectra (Groenewegen et al. 2007). The C₂H₂ molecular bands are included in the model for the stellar atmosphere, which is used as an input to DUSTY models, but the observed bands are stronger than predicted. As discussed later, the parameters used in the models are appropriate for Galactic stars, and the metallicity dependence of the molecular band strengths may well contribute to this problem. There could be additional effects, such as some C₂H₂ originating from the extended atmosphere which is lifted by the pulsation above the photosphere (e.g. Hron et al. 1998; Ohnaka et al. 2007). Circumstellar molecular absorption (e.g. Aoki, Tsuji & Ohnaka 1999; Matsuura et al. 2006) is also not included in DUSTY.

The mass-loss rates for the three reddest stars are of the order of $5 \times 10^{-6} \,\text{M}_{\odot} \,\text{yr}^{-1}$. This would be only a moderate mass-loss rate for Galactic carbon stars. The comparison assumes that the gas-to-dust ratio in the Fornax dSph galaxy is the same as in our Galaxy.

The bolometric luminosities estimated from fitting a radiative transfer model are comparable to other estimates, such as those from near-IR, J - K, colours (Lagadec et al. 2007b). The values range between log $L[L_{\odot}] = 3.9$ and 4.1, which is comparable to the mean luminosities found for carbon stars in the LMC, but slightly higher than those found for SMC stars (Buchanan et al. 2006; Zijlstra et al. 2006; Lagadec et al. 2007a).

Fig. 9 shows the derived mass-loss rate as a function of luminosity and a comparison with values for LMC and SMC stars (Groenewegen et al. 2007). Little metallicity dependence of the mass-loss rate



Figure 9. Mass-loss rate $(M_{\odot} yr^{-1})$ as a function of luminosity (L_{\odot}) for our sample and for the LMC and SMC samples (Groenewegen et al. 2007). Fornax stars show high mass-loss rates for their luminosities. The dotted line is the fit to the luminosity versus mass-loss rate relation for Galactic carbon stars (Groenewegen et al. 1998).

is found, at least for the dust mass-loss rate. The Fornax mass-loss rates are at the upper end of the SMC mass-loss rates at a given luminosity. However, we are unable to fit the Fornax stars with lower mass-loss rates (Fornax 4-25 and 6-13), so there is an obvious selection effect. LMC stars appear to reach a higher mass-loss rate than the SMC and Fornax stars at a given luminosity, as noted by Groenewegen et al. (2007).

The assumptions made in deriving the total mass-loss rates are the expansion velocity of the circumstellar shell and the conversion factor from dust mass-loss rate to gas mass-loss rate.

Groenewegen et al. (2007) assumed an expansion velocity of 10 km s^{-1} for all the stars in their study. The velocity is derived from DUSTY fitting in our work and we find that the Fornax stars have expansion velocities in the 16 to 25 km s^{-1} range. This gives us larger mass-loss rates for the Fornax stars by a factor of up to 2.5 over those obtained from the Groenewegen et al. (2007) assumption. Nevertheless, in the log scale of Fig. 9, this is a minor effect.

The gas-to-dust mass ratio is assumed to be 200 in both this work and by Groenewegen et al. (2007). This ratio may have a metallicity dependence, and van Loon (2000) estimates $\log \psi \propto -1.0$ [Z/H] for carbon stars, where ψ is the gas-to-dust ratio. If this is correct, the gas-to-dust ratio in Fornax will be about twice what it is in the SMC and the Fornax stars will have higher gas mass-loss rates than derived above. We discuss this further below (Section 6.2).

Fig. 10 shows the mass-loss rate as a function of IR colour. The wavelength bands are defined so as to avoid major molecular or dust features, thus the colour measures the 'continuum' spectrum reasonably well. In Fig. 10, a correlation is found between the IR colour and the mass-loss rate. As mentioned in Groenewegen et al. (2007), this colour gives the approximate mass-loss rate with a factor of 2 to 5 uncertainty, depending on the colour.

5.3 A comparison with IR colour method

Lagadec et al. (2007b) estimated the dust mass-loss rate of two of the sources in our sample, using the IR colour index K - [11].



Figure 10. Mass-loss rate as a function of IR colour [6.4]-[9.3].

Their dust mass-loss rates have been converted to gas mass-loss rates assuming a gas-to-dust ratio of 200 and are listed in Table 4; they are consistent with our estimates to within a factor of 2. At least some of the difference will be due to the input stellar spectra, i.e., our work uses a hydrostatic model while Lagadec et al. (2007b) assume a BB. Nevertheless, the consistency shows that the IR colour method can be used to estimate the mass-loss rates, as found for Galactic AGB stars (Whitelock et al. 1994; Le Bertre 1997).

The mid-IR colour also shows a correlation with mass-loss rate (Fig. 10), especially when using colours that minimize the influence of strong molecular bands. Note that unlike some colour measurements, e.g. K - [12], our values are constructed from spectra and are therefore simultaneous and not affected by variability.

6 DISCUSSION

6.1 Luminosity of carbon stars

The carbon stars in the Fornax dSph galaxy tend to have fainter absolute *K* magnitudes at a given J - K colour (Fig. 1) than LMC carbon stars. Demers et al. (2002) pointed out that the mean magnitudes of carbon-rich stars at the tip of the AGB are $\langle M_K \rangle = -7.91$ and -7.88 mag for the LMC and SMC, respectively, while the mean magnitude of 26 carbon stars in Fornax gives $\langle M_K \rangle = -7.68$ mag. The difference between the LMC and SMC samples was also discussed in Lagadec et al. (2007a, fig. 2). Demers et al. (2002) concluded that the low $\langle M_K \rangle$ is due to the lack of younger (0.3–1 Gyr) carbon stars in Fornax. The population analysis by Battaglia et al. (2006) also supports this interpretation; the star formation rate declined during the period from a few hundred Gyr to 1 Gyr ago, so we are unlikely to find young (0.3–1 Gyr) carbon stars in Fornax. In other words, AGB stars in the Fornax dSph Galaxy are, on average, probably older than those found in the LMC and the SMC.

6.2 Metallicity dependence of mass-loss rates

The ratio of SiC to continuum is smaller in Fornax stars than in our Galaxy or in the LMC, but comparable to that in the SMC. The Fornax spectra are well fitted if the mixture of dust grains is 5 per cent SiC and 95 per cent amorphous carbon, although the error in our estimate is large. As Groenewegen et al. (2007) found, the fraction of SiC dust grains is lower in carbon stars in the LMC and SMC than those in our Galaxy (10 per cent; e.g. Groenewegen 1995). Stars in the Fornax dSph galaxy follow this same trend with metallicity.

The fraction of SiC dust in the grains is probably correlated with the silicon elemental abundance in the atmospheres of carbon stars. The temperature in the helium burning shell does not become high enough for α -captures on magnesium, and the abundance of ²⁸Si can be assumed to be unchanged during the AGB phase. The abundance of silicon has been measured in two red giants in the Fornax dSph galaxy (Shetrone et al. 2003). They find silicon abundances in metalpoor stars higher than implied by a simple scaling from the metallicity as represented by [Fe/H]. These two stars have approximately 5 and 25 per cent of the solar silicon abundance ([Si/H] = -1.3 and -0.6), while their iron abundances are [Fe/H] = -1.60 and -0.67. Thus, the metal-poor star shows a silicon abundance excess, presumably due to the detailed history of its formation (Shetrone et al. 2003).

A mass fraction of 5 per cent SiC grains and a gas-to-dust ratio of 200 in mass, implies that 25 per cent of the silicon atoms are in the dust, if the envelope has a solar metallicity. Here we assume the solar abundance of silicate is 7.55, on a scale of $\log N(H) =$ 12. For [Si/H] = -0.6, all silicon atoms would need to be in the SiC dust grains. We expect our stars to have somewhat lower metallicity, corresponding to the peak of the stellar metallicity distribution, say $[Fe/H] \sim -1.0$ and the silicon abundance ([Si/H]) could be ~ -0.9 . Inverting the estimation, the gas-to-dust ratio needs to be 400 or larger, to account for the SiC mass fraction of 5 per cent. If we consider the uncertainty in this estimate of the SiC fraction, then the gas-to-dust ratio must be in the range 800 to 240 to provide a fraction of 10 to 3 per cent of the dust as SiC. This may be the first, albeit indirect, indication for a higher gas-to-dust ratio at significantly subsolar metallicity. If it is correct, then the mass-loss rates in AGB stars at these low metallicities could be even higher than values given in Table 4. The uncertainty in the gas-to-dust ratio discussed here largely depends on the uncertainty of our estimated SiC fraction. Various parameters, but particularly the optical constants of the dust grains and the dust grain size distribution, require refinement if further progress is to be made. Furthermore, the exact compositions mixed in with the SiC dust grains (Heck et al. 2007) might change towards lower metallicity, resulting in different SiC dust masses. Although it is possible that the gas-to-dust ratio increases at lower metallicity the actual values remain very uncertain. Because of these uncertainties, we continue to assume that the gas-to-dust ratio is 200 for the remaining discussion.

The dust mass is dominated by amorphous carbon. The quantity of amorphous carbon grains is affected by two factors, viz. C/O ratio (Habing, Tignon & Tielens 1994) and initial metallicity (Zuckerman & Dyck 1989; van Loon 2000; Wahlin et al. 2006). The lack of evidence for lower mass-loss rates at lower metallicity is explained with a higher C/O ratio towards low metallicity. A strong dependence of mass-loss rate on C/O ratio is reported from hydrodynamical models for carbon-rich stars (Mattsson, Wahlin & Höfner 2007). This is a consequence of the stability of the CO molecule; in carbon stars all available oxygen atoms are locked into carbon monoxide and excess carbon atoms are available to form carbon-bearing molecules and amorphous carbon dust grains. As new carbon atoms reach the surface of these AGB stars, the C/O ratio increases and the amount of carbon available to form grains also increases. The C/O abundance of the target stars in the LMC, the SMC and Fornax have not been measured directly. A number of carbon stars in

the SMC have C/O ratios of 1.05 to 1.1 (de Laverny et al. 2006). That sample probably contains blue carbon stars, which have little or no circumstellar envelope. The C/O ratio increases towards the end of the AGB phase, but such stars develop a thick circumstellar envelope and hydrostatic models cannot be used for them, because the molecular lines are filled in by emission from circumstellar dust grains. C/O ratios above 2 are found in LMC and SMC planetary nebulae (Leisy & Dennefeld 1996; Stanghellini, Shaw & Gilmore 2005). At metallicities below those of the SMC and LMC, the final C/O ratios are expected to be even higher due to low initial oxygen abundance, and an additional effect of efficient third-dredge up (Vassiliadis & Wood 1993; Marigo, Girardi & Chiosi 2003; Izzard et al. 2004). The C/O ratios of our targets are therefore likely to significantly exceed 1.1, leading to the observed efficient amorphous dust production.

This result, i.e., the lack of dependence of mass-loss rates on metallicity for carbon-rich stars, is in contrast to what is found for oxygen-rich stars, where there is an obvious metallicity dependence (Wood et al. 1992). This is because dust-driven winds in oxygen-rich stars are dominated by the amount of silicate, and ultimately by the intrinsic silicon elemental abundance (Bowen & Willson 1991; Willson 2006). The expansion velocities of oxygenrich shells are also smaller at low metallicities (Wood et al. 1992; Zijlstra et al. 1996; Marshall et al. 2004). In contrast, carbon stars are strongly affected by carbon production during the AGB phase; thus the different dependence of mass-loss rate on metallicity can be understood.

6.3 Metallicity dependence of molecular bands

The molecular band strengths of carbon-rich stars show a metallicity dependence (Cohen et al. 1981; van Loon, Zijlstra & Groenewegen 1999; Matsuura et al. 2002). Recent studies using infrared spectrometers on the Very Large Telescope (Matsuura et al. 2005; van Loon et al. 2006) showed that the C_2H_2 equivalent widths increase towards lower metallicity, based on samples from the Galaxy, the LMC and the SMC.

The carbon-rich stars in the Fornax dSph galaxy follow the same trend of as those found in similar SMC stars. For IR colours [6.4]– [9.3] > 0.5, the largest C_2H_2 equivalent widths are found for stars in the SMC and the Fornax dSph galaxy. Although Fornax has a lower metallicity than the SMC, no further metallicity dependence is seen below the SMC value.

As discussed in the previous section, a higher C/O ratio is likely in carbon-rich stars at lower metallicity. C_2H_2 formation relies on excess carbon atoms after all oxygen atoms are locked into carbon monoxide. Thus the higher C/O ratio results in the higher abundance of C_2H_2 at lower metallicity (Matsuura et al. 2005). Fig. 13 in Matsuura et al. (2005) shows that the fraction of C_2H_2 in the atmosphere increases drastically up to C/O ~1.6, but at ratios higher than this the amount of C_2H_2 increases only slowly. This is because C_2H_2 is reactive with carbon resulting in increasingly larger carbon bearing molecules. Thus, below the SMC metallicity, little metallicity dependence of the C_2H_2 equivalent width is found, because this molecular abundance becomes insensitive to the C/O ratio.

 C_2H_2 is thought to be a parent molecule in the formation of polycyclic aromatic hydrocarbons (PAHs), as indicated by chemical models (Allamandola, Tielens & Barker 1989). It is still unknown whether PAHs are formed during the AGB phase or afterwards. They may already form in the atmospheres or inner circumstellar envelopes of carbon-rich AGB stars, where the gas density is reasonably high and chemical reactions occur easily (Frenklach & Feigelson 1989). The non-detection of PAHs from AGB stars might be due to a lack of ultraviolet radiation, which is probably required to excite PAHs. PAHs might be photon processed during post-AGB evolution (Sloan et al. 2007). Alternatively, PAHs may be formed during the post-AGB or protoplanetary nebula phase, where PAHs are commonly found. The chemistry in this phase is dominated by strong radiation from the central star (Woods et al. 2002, 2003). If PAHs are indeed formed during the AGB phase, an overabundance of C_2H_2 in a low metal environment will affect the growth of these important molecules.

The overabundance of C_2H_2 at low metallicity will remain through the post-AGB phase, as is illustrated by the fact that the LMC post-AGB star SMP 11 has more abundant carbon-bearing molecules (Bernard-Salas et al. 2006) than comparable Galactic post-AGB stars (Cernicharo et al. 2001). Reyniers et al. (2007) suggested a lower limit of C/O = 2 for one LMC post-AGB star. This overabundance of C_2H_2 may also occur in an SMC post-AGB star (Kraemer et al. 2006). Thus, at subsolar metallicity, it is likely that C_2H_2 molecules are abundant in carbon-rich evolved stars.

6.4 Gas supply from AGB stars to the ISM of the Fornax dSph galaxy

From near-IR colours, Gullieuszik et al. (2007) found about 100 AGB candidates in the centre $(18.5 \times 18.5 \text{ arcmin}^{-1})$ of the Fornax dSph galaxy. Their IR colours suggest that at least 20 of these AGB stars are carbon rich (J - K > 1.4 and K < 14 mag) (e.g. Cioni & Habing 2003). The reddest carbon stars found by Gullieuszik et al. (2007) have J - K = 2.4 mag. An estimate of mass-loss rate can be obtained from the J - K colours of these 20 AGB stars of the order of $2 \times 10^{-7} \,\mathrm{M_{\odot}} \,\mathrm{yr^{-1}}$ or less, although this number is not very precise owing to the large dispersion of the J - K versus mass-loss rate relation found by Le Bertre (1997). Including our carbon stars, the total mass-loss rate from known carbon-rich stars in Fornax dSph galaxy is estimated to be about $2 \times 10^{-5} \, M_{\odot} \, yr^{-1}$ at most, adopting a gas-to-dust ratio of 200. Three stars in our sample (Fornax 13-23, 12-4 and 3-129) contribute more than two-thirds of the total massloss rate. Studies of Galactic AGB stars showed that stars with higher mass-loss rates (above $1 \times 10^{-6} \,\mathrm{M_{\odot} yr^{-1}}$) dominate the total gas output from AGB stars (Le Bertre et al. 2001). Therefore, the much bluer AGB stars in Fornax are assumed to contribute very little to the total mass lost from the entire AGB population.

This total mass-loss rate is smaller than the value found for Wolf Lundmark Melotte (WLM), a dwarf irregular galaxy in the Local Group with a similar metallicity range to Fornax. Jackson et al. (2007) estimated the total mass-loss rate from AGB stars as [0.7-2.4] $\times 10^{-3}$ M_{\odot} yr⁻¹, which is a factor of 50 larger in WLM than the Fornax dSph galaxy. The number ratio of carbon- to oxygen-rich stars is reported to be 12 in WLM (Battinelli, Demers & Letarte 2003); thus the mass-loss rate from carbon-rich stars is almost identical to that from the entire AGB population. The visual absolute magnitudes (M_V) of the galaxies are -14.4 and -13.1 for WLM and Fornax, respectively. If M_V is a measure of the number of stars in a galaxy, WLM has a two to three times larger the number of stars than Fornax has; the number of stars in the galaxies is insufficient to explain the difference in total mass-loss rate from carbon-rich stars. The assumed gas-to-dust ratio is a factor of 6 to 13 larger in Jackson et al. (2007), but still not sufficient to explain the result. The difference might be caused by the presence of younger stars in WLM, and the absence of such stars in Fornax. In WLM, a recent increase in star formation rate is indicated starting 1–2.5 Gyr ago (Hodge & Miller 1995). Young (<1 Gyr) carbon stars dominate the mass output from AGB stars (e.g. Le Bertre et al. 2001), and this population is missing in the Fornax dSph galaxy.

As a dwarf irregular galaxy, WLM has a much larger H_I mass than the Fornax dSph: 40 per cent of its total mass compared to <0.1 per cent for Fornax (Mateo 1998). A removal of gas from dSph in early phase is a consequence of this difference (Grebel, Gallagher & Harbeck 2003). However, the feedback from the young AGB stars also accentuates the difference in gaseous contents, on top of the influence of the galaxy's initial gas content.

6.5 Period of Fornax 4-15

The absolute magnitude of Fornax 4-25 is estimated as $M_{bol} = -4.09$ mag, using the *K* magnitude and colour-dependent bolometric correction (Whitelock et al. 2006). The period–luminosity relation for Galactic carbon stars (Whitelock et al. 2006) gives $\log P = 2.3$, where *P* is the period in days, for this luminosity. The observed period of Fornax 4-15 is $\log P = 2.5$ (Demers & Irwin 1987), thus this Fornax star fits on the period–luminosity relation for Galactic stars very well. No obvious metallicity dependence has been found for the AGB period–luminosity relation. This period and mass-loss rate relation for galactic star gives mass-loss rate of $10^{-6} \,\mathrm{M_{\odot} yr^{-1}}$ at $\log P = 2.5$. This mass-loss rate is consistent with the upper limit (<1.3 × $10^{-6} \,\mathrm{M_{\odot} yr^{-1}}$) obtained by this work.

7 CONCLUSIONS

The 5–14 µm spectra of AGB stars in the Fornax dSph galaxy confirm the prevalence of carbon-rich stars at low metallicity ([Fe/H] ~ -1). Stars with J - K > 3 show spectra dominated by C₂H₂ molecules, together with CO bands and SiC dust excess, while only a trace of the 7.5 µm C₂H₂ is found in stars with J - K < 3. Detection of the SiC dust excess shows that these AGB stars have developed circumstellar envelopes through mass loss.

We compare our results for Fornax with carbon-rich stars in our own Galaxy, the LMC and the SMC. The Fornax dSph galaxy is the lowest metallicity galaxy for which the carbon-rich star population has been studied in the mid-IR. The C_2H_2 molecular bands, with respect to the IR colour, tend to be stronger towards lower metallicity. No reduction in the mass-loss rate is found for lower metallicity carbon stars. The dust composition does depend on the metallicity, with a lower mass fraction of SiC dust grains, but the amount of amorphous carbon, which constitutes the dominant fraction of dust grains, seem to be unchanged. If the fraction of SiC dust grain is correct, the gas-to-dust ratio could be as high as 400 at one tenth of the solar metallicity, but a more detailed investigation of gas mass-loss rates is required. This study of AGB mass loss in a low-metallicity environment should be broadly useful for other studies of dust and gas recycling within metal-poor galaxies.

Our study reveals that the gas supply from AGB stars into the ISM is not lower at lower metallicity, but the fraction of SiC dust grains does appear to be lower. The total amount of gas expelled from AGB stars is less in the Fornax dSph galaxy than in the dwarf irregular WLM, in spite of a similar total mass and metallicity. The difference is attributed to the younger stellar population in WLM.

ACKNOWLEDGMENTS

MM thanks Prof. Arimoto for discussion about populations in Fornax dSph galaxy. MM is a JSPS fellow.

REFERENCES

- Aaronson M., Mould J., 1980, ApJ, 240, 804
- Allamandola L. J., Tielens G. G. M., Barker J. R., 1989, ApJS, 71, 733
- Anders E., Grevesse N., 1989, Geochim. Cosmochim. Acta, 53, 197
- Aoki W., Tsuji T., Ohnaka K., 1999, A&A, 350, 945
- Azzopardi M., Muratorio G., Breysacher J., Westerlund B. E., 1999, in Whitelock P., Cannon R., eds, Proc. IAU Symp. 192, The Stellar Content of Local Group Galaxies. Astron. Soc. Pac., San Francisco, p. 144 Battaglia G. et al., 2006, A&A, 459, 423
- Battinelli P., Demers S., Letarte B., 2003, AJ, 125, 1298
- Bernard-Salas J., Peeters E., Sloan G. C., Cami J., Guiles S., Houck J. R., 2006, ApJ, 652, L29
- Bertelli G., Bressan A., Chiosi C., Fagotto F., Nasi E., 1994, A&AS, 106, 275
- Bouchard A., Carignan C., Staveley-Smith L., 2006, AJ, 131, 2913
- Bowen G. H., Willson L. A., 1991, ApJ, 375, L53
- Buchanan C. L., Kastner J. H., Forrest W. J., Hrivnak B. J., Sahai R., Egan M., Frank A., Barnbaum C., 2006, AJ, 132, 1890
- Busso M., Guandalini R., Persi P., Corcione L., Ferrari-Toniolo M., 2007, AJ, 133, 2310
- Cernicharo J., Heras A. M., Pardo J. R., Tielens A. G. G. M., Guélin M., Dartois E., Neri R., Waters L. B. F. M., 2001, ApJ, 546, L127
- Cioni M.-R. L., Habing H. J., 2003, A&A, 402, 133
- Cioni M.-R. L., Marquette J.-B., Loup C., Azzopardi M., Habing H. J., Lasserre T., Lesquoy E., 2001, A&A, 377, 945
- Cohen J. G., Persson S. E., Elias J. H., Frogel J. A., 1981, ApJ, 249, 481
- Cohen M., Wheaton Wm. A., Megeath S. T., 2003, AJ, 126, 1090
- Cole A. A., Tolstoy E., Gallagher J. S., III, Smecker-Hane T. A., 2005, AJ, 129, 1465
- de Laverny P., Abia C., Domínguez I., Plez B., Straniero O., Wahlin R., Eriksson K., Jorgensen U. G., 2006, A&A, 446, 1107
- Demers S., Irwin M. J., 1987, MNRAS, 226, 943
- Demers S., Kunkel W. E., 1979, PASP, 91, 761
- Demers S., Grondin L., Kunkel W. E., 1990, PASP, 102, 632
- Demers S., Dallaire M., Battinelli P., 2002, AJ, 123, 3428
- Dwek E., 1998, ApJ, 501, 643
- Frenklach M., Feigelson E. D., 1989, ApJ, 341, 372
- Frogel J. A., Blanco V. M., Cohen J. G., McCarthy M. F., 1982, ApJ, 252, 133
- Glass I. S., Nagata T., 2000, Mon. Notes Astron. Soc. South Afr., 59, 110
- Grebel E. K., Gallagher J. S., III, Harbeck D., 2003, AJ, 125, 1926
- Groenewegen M. A. T., 1995, A&A, 293, 463
- Groenewegen M. A. T., Whitelock P. A., Smith C. H., Kerschbaum F., 1998, MNRAS, 293, 18
- Groenewegen M. A. T. et al., 2007, MNRAS, 376, 313
- Gullieuszik M., Held E. V., Rizzi L., Saviane I., Momany Y., Ortolani S., 2007, A&A, 467, 1025
- Habing H. J., Tignon J., Tielens A. G. G. M., 1994, A&A, 286, 523
- Hanner M. S., 1988, NASA Conf. Publ., 3004, 22
- Heck P. R., Marhas K. K., Hoppe P., Gallino R., Baur H., Wieler R., 2007, ApJ, 656, 1208
- Hodge P., Miller B. W., 1995, ApJ, 451, 176
- Houck J. R. et al., 2004, ApJS, 154, 18
- Hron J., Loidl R., Höfner S., Jorgensen U. G., Aringer B., Kerschbaum F., 1998, A&A, 335, L69
- Irwin M., Hatzidimitriou D., 1995, MNRAS, 277, 1354
- Ivezić Z., Elitzur M., 1995, ApJ, 445, 415
- Ivezić Z., Elitzur M., 1997, MNRAS, 287, 799
- Izzard R. G., Tout C. A., Karakas A. I., Pols O. R., 2004, MNRAS, 350, 407
- Jackson D. C., Skillman E. D., Gehrz R. D., Polomski E., Woodward C. E., 2007, ApJ, 656, 818
- Jørgensen U. G., Hron J., Loidl R., 2000, A&A, 356, 253
- Kniazev A. Y., Grebel E. K., Pustilnik S. A., Pramskij A. G., 2007, A&A, 468, 121
- Kraemer K. E., Sloan G. C., Bernard-Salas J., Price S. D., Egan M. P., Wood P. R., 2006, ApJ, 652, L25

Lagadec E. et al., 2007a, MNRAS, 376, 1270

- Lagadec E. et al. 2007b, MNRAS, in press (doi:10.1111/j.1365-2966. 2007.12561.x)
- Lattanzio J., Forestini M., 1999, in Le Bertre T., Lèbre A., Waelkens C., eds, IAU Symp. 191, Asymptotic Giant Branch Stars. Astron. Soc. Pac., San Francisco, p. 31
- Le Bertre T., 1997, A&A, 324, 1059
- Le Bertre T., Matsuura M., Winters J. M., Murakami H., Yamamura I., Freund M., Tanaka M., 2001, A&A, 376, 997
- Leisy P., Dennefeld M., 1996, A&AS, 116, 95
- Loidl R., Lançon A., Jorgensen U. G., 2001, A&A, 371, 1065
- Maeder A., 1992, A&A, 264, 105
- Marigo P., Girardi L., Chiosi C., 2003, A&A, 403, 225
- Marshall J. R., van Loon J. Th., Matsuura M., Wood P. R., Zijlstra A. A., Whitelock P. A., 2004, MNRAS, 355, 1348
- Mateo M. L., 1998, ARA&A, 36, 435
- Mathis J. S., Rumpl W., Nordsieck K. H., 1977, ApJ, 217, 425
- Matsuura M., Zijlstra A. A., van Loon J. Th., Yamamura I., Markwick A. J., Woods P. M., Waters L. B. F. M., 2002, ApJ, 580, L133
- Matsuura M. et al., 2005, A&A, 434, 691
- Matsuura M. et al., 2006, MNRAS, 371, 415
- Mattsson L., Wahlin R., Höfner S., 2007, in Vazdekis A., Peletier R. F., eds, Proc. IAU Symp. 241, Stellar Populations as Building Blocks of Galaxies. Cambridge Univ. Press, Cambridge, p. 37
- Menzies J., Feast M., Tanabé T., Whitelock P., Nakada Y., 2002, MNRAS, 335, 923
- Ohnaka K., Driebe T., Weigelt G., Wittkowski M., 2007, A&A, 466, 1099
- Pégourié B., 1988, A&A, 194, 335
- Pont F., Zinn R., Gallart C., Hardy E., Winnick R., 2004, AJ, 127, 840
- Reyniers M., Abia C., van Winckel H., Lloyd Evans T., Decin L., Eriksson K., Pollard K. R., 2007, A&A, 461, 641
- Saviane I., Held E. V., Bertelli G., 2000, A&A, 355, 56
- Shetrone M., Venn K. A., Tolstoy E., Primas F., Hill V., Kaufer A., 2003, AJ, 125, 684
- Skrutskie M. F. et al., 2006, AJ, 131, 1163
- Sloan G. C., Kraemer K. E., Price S. D., Shipman R. F., 2003, ApJS, 147, 379
- Sloan G. C. et al., 2006, ApJ, 645, 1118
- Sloan G. C. et al., 2007, ApJ, 664, 1144
- Smith V. V., Lambert D. L., 1989, ApJ, 345, L75
- Stanghellini L., Shaw R. A., Gilmore D., 2005, ApJ, 622, 294
- Strader J., Brodie J. P., Forbes D. A., Beasley M. A., Huchra J. P., 2003, AJ, 125, 1291
- Thompson G. D., Corman A. B., Speck A. K., Dijkstra C., 2006, ApJ, 652, 1654
- Tolstoy E., Irwin M. J., Cole A. A., Pasquini L., Gilmozzi R., Gallagher J. S., 2001, MNRAS, 327, 918
- Tolstoy E., Venn K. A., Shetrone M., Primas F., Hill V., Kaufer A., Szeifert T., 2003, AJ, 125, 707
- van den Bergh S., 1999, Cambridge Astrophys. Ser. Vol. 35, The Galaxies of the Local Group. Cambridge Univ. Press, Cambridge
- van Loon J. Th., 2000, A&A, 354, 125
- van Loon J. Th., Zijlstra A. A., Groenewegen M. A. T., 1999, A&A, 346, 805
- van Loon J. Th., Marshall J. R., Matsuura M., Zijlstra A. A., 2003, MNRAS, 341, 1205
- van Loon J. Th., Marshall J. R., Cohen M., Matsuura M., Wood P. R., Yamamura I., Zijlstra A. A., 2006, A&A, 447, 971
- Vassiliadis E., Wood P. R., 1993, ApJ, 413, 641
- Wahlin R., Eriksson K., Gustafsson B., Hinkle K. H., Lambert D. L., Ryde N., Westerlund B., 2006, Mem. Soc. Astron. Ital., 77, 955
- Werner M. W. et al., 2004, ApJS, 154, 1
- Westerlund B. E., Edvardsson B., Lundgren K., 1987, A&A, 178, 41
- Whitelock P., Menzies J., Feast M., Marang F., Carter B., Roberts G., Catchpole R., Chapman J., 1994, MNRAS, 267, 711
- Whitelock P. A., Feast M. W., Marang F., Groenewegen M. A. T., 2006, MNRAS, 369, 751

- Willson L. A., 2006, in Stanghellini L., Walsh J. R., Douglas N. G., eds, Proc. ESO, Planetary Nebulae Beyond the Milky Way. Springer-Verlag, Berlin, p. 99
- Wood P. R., Whiteoak J. B., Hughes S. M. G., Bessell M. S., Gardner F. F., Hyland A. R., 1992, ApJ, 397, 552
- Wood P. R. et al., 2007, Kerschbaum F., Charbonnel C., eds, ASP Conf. Ser. Vol. 378, Why Galaxies Care About AGB stars. Astron. Soc. Pac., San Francisco, in press
- Woods P. M., Millar T. J., Zijlstra A. A., Herbst E., 2002, ApJ, 574, L167
- Woods P. M., Millar T. J., Herbst E., Zijlstra A. A., 2003, A&A, 402, 189
- Yamamura I., de Jong T., Justtanont K., Cami J., Waters L. B. F. M., 1997, Ap&SS, 255, 351
- Young L. M., 1999, AJ, 117, 1758
- Zijlstra A. A., Loup C., Waters L. B. F. M., Whitelock P. A., van Loon J. Th., Guglielmo F., 1996, MNRAS, 279, 32
- Zijlstra A. A. et al., 2006, MNRAS, 370, 1961
- Zuckerman B., Dyck H. M., 1989, A&A, 209, 119

This paper has been typeset from a $T_{\ensuremath{E}} X/ I \!\! \ensuremath{\Delta} T_{\ensuremath{E}} X$ file prepared by the author.